



A Level Biology



Transition Induction Pack

Contents

How to use this book
Kerboodle

	viii	xi
Section 1		
Biological molecules		
1 Biological molecules	2	84
1.1 Introduction to biological molecules	4	84
1.2 Carbohydrates and monosaccharides	4	87
1.3 Carbohydrates – disaccharides and polysaccharides	8	89
1.4 Starch, glycogen and cellulose	10	93
1.5 Lipids	13	95
1.6 Proteins	16	99
1.7 Enzyme action	19	102
1.8 Factors affecting enzyme action	23	102
1.9 Enzyme inhibition	26	104
Practice questions	32	106
2 Nucleic acids	34	109
2.1 Structure of RNA and DNA	36	111
2.2 DNA replication	36	115
2.3 Energy and ATP	42	119
2.4 Water and its functions	46	122
Practice questions	50	124
Section 1 summary	52	126
Section 1 practice questions	54	
Section 2		
Cells	56	128
3 Cell structure	58	130
3.1 Methods of studying cells	58	130
3.2 The electron microscope	61	133
3.3 Microscopic measurements and calculations	64	135
3.4 Eukaryotic cell structure	67	137
3.5 Cell specialisation and organisation	73	139
3.6 Prokaryotic cells and viruses	75	142
3.7 Mitosis	77	144
3.8 The cell cycle	80	146
Practice questions	82	151
Section 2 summary	84	155
Section 2 practice questions	86	158
Section 3		
Organisms exchange substances with their environment	128	
6 Exchange	130	
6.1 Exchange between organisms and their environment	130	
6.2 Gas exchange in single-celled organisms and insects	133	
6.3 Gas exchange in fish	135	
6.4 Gas exchange in the leaf of a plant	137	
6.5 Limiting water loss	139	
6.6 Structure of the human gas-exchange system	142	
6.7 The mechanism of breathing	144	
6.8 Exchange of gases in the lungs	146	
6.9 Enzymes and digestion	151	
6.10 Absorption of the products of digestion	155	
Practice questions	158	
Section 4		
Transport across cell membranes	84	
4.1 Structure of the cell surface membrane	84	
4.2 Diffusion	87	
4.3 Osmosis	89	
4.4 Active transport	93	
4.5 Co-transport and absorption of glucose in the ileum	95	
Practice questions	99	
5 Cell recognition and the immune system	102	
5.1 Defence mechanisms	102	
5.2 Phagocytosis	104	
5.3 T lymphocytes and cell mediated immunity	106	
5.4 B lymphocytes and humoral immunity	109	
5.5 Antibodies	111	
5.6 Vaccination	115	
5.7 Human immunodeficiency virus (HIV)	119	
Practice questions	122	
Section 2 summary	124	
Section 2 practice questions	126	
Section 3		
Organisms exchange substances with their environment	128	
6 Exchange	130	
6.1 Exchange between organisms and their environment	130	
6.2 Gas exchange in single-celled organisms and insects	133	
6.3 Gas exchange in fish	135	
6.4 Gas exchange in the leaf of a plant	137	
6.5 Limiting water loss	139	
6.6 Structure of the human gas-exchange system	142	
6.7 The mechanism of breathing	144	
6.8 Exchange of gases in the lungs	146	
6.9 Enzymes and digestion	151	
6.10 Absorption of the products of digestion	155	
Practice questions	158	
Section 4		
Mass transport	161	
7.1 Haemoglobin	161	
7.2 Transport of oxygen by haemoglobin	163	
7.3 Circulatory system of a mammal	168	
7.4 The structure of the heart	170	
7.5 The cardiac cycle	174	
7.6 Blood vessels and their functions	178	
7.7 Transport of water in the xylem	183	
7.8 Transport of organic molecules in the phloem	188	
7.9 Investigating transport in plants	191	
Practice questions	194	
Section 3 summary	196	
Section 3 practice questions	198	
Section 4		
Genetic information, variation and relationships between organisms	200	
8 DNA, genes and protein synthesis	202	
8.1 Genes and the triplet code	202	
8.2 DNA and chromosomes	205	
8.3 The structure of ribonucleic acid	208	
8.4 Protein synthesis – transcription and splicing	211	
8.5 Protein synthesis – translation	213	
Practice questions	217	
9 Genetic diversity	220	
9.1 Mutations	220	
9.2 Meiosis and genetic variation	224	
9.3 Genetic diversity and adaptation	229	
9.4 Types of selection	231	
Practice questions	235	
10 Biodiversity	237	
10.1 Species and taxonomy	237	
10.2 Diversity within a community	243	
10.3 Species diversity and human activity	246	
10.4 Investigating diversity	249	
10.5 Quantitative investigations of variation	253	
Practice questions	257	
Section 4 summary	260	
Section 4 practice questions	262	
Section 5		
Energy transfer in and between organisms	266	
11 Photosynthesis	268	
11.1 Overview of photosynthesis	268	
11.2 The light-dependent reaction	271	
11.3 The light-independent reaction	275	
Practice questions	281	
12 Respiration	283	
12.1 Glycolysis	283	
12.2 Link reaction and Krebs cycle	286	
12.3 Oxidative phosphorylation	289	
12.4 Anaerobic respiration	293	
Practice questions	296	
13 Energy and ecosystems	298	
13.1 Food chains and energy transfer	298	
13.2 Energy transfer and productivity	300	
13.3 Nutrient cycles	306	
13.4 Use of natural and artificial fertilisers	311	
13.5 Environmental issues concerning use of nitrogen-containing fertilisers	313	
Practice questions	315	
Section 5 summary	318	
Section 5 practice questions	320	
Section 6		
Organisms respond to changes in their environments	324	
14 Response to stimuli	326	
14.1 Survival and response	326	
14.2 Plant growth factors	328	
14.3 A reflex arc	334	
14.4 Receptors	337	
14.5 Control of heart rate	340	
Practice questions	344	

Assessments

marksphysicshelp MPH

Paper 1	+	Paper 2	+	Paper 3
What's assessed Any content from topics 1–4, including relevant practical skills		What's assessed Any content from topics 5–8, including relevant practical skills		What's assessed Any content from topics 1–8, including relevant practical skills
Assessed - written exam: 2 hours 91 marks 35% of A-level		Assessed - written exam: 2 hours 91 marks 35% of A-level		Assessed - written exam: 2 hours 78 marks 30% of A-level
Questions 76 marks: a mixture of short and long answer questions 15 marks: extended response questions		Questions 76 marks: a mixture of short and long answer questions 15 marks: comprehension question		Questions 38 marks: structured questions, including practical techniques 15 marks: critical analysis of given experimental data 25 marks: one essay from a choice of two titles

8.2 A-level required practical activities

marksphysicshelp MPH

The following practicals must be carried out by all students taking this course. Written papers will assess knowledge and understanding of these, and the skills exemplified within each practical.

Required activity	Apparatus and technique reference
1. Investigation into the effect of a named variable on the rate of an enzyme-controlled reaction	a, b, c, f, l
2. Preparation of stained squashes of cells from plant root tips; set-up and use of an optical microscope to identify the stages of mitosis in these stained squashes and calculation of a mitotic index	d, e, f
3. Production of a dilution series of a solute to produce a calibration curve with which to identify the water potential of plant tissue	c, h, j, l
4. Investigation into the effect of a named variable on the permeability of cell-surface membranes	a, b, c, j, l
5. Dissection of animal or plant gas exchange or mass transport system or of organ within such a system	e, h, j
6. Use of aseptic techniques to investigate the effect of antimicrobial substances on microbial growth	c, i
7. Use of chromatography to investigate the pigments isolated from leaves of different plants, eg leaves from shade-tolerant and shade-intolerant plants or leaves of different colours	b, c, g
8. Investigation into the effect of a named factor on the rate of dehydrogenase activity in extracts of chloroplasts	a, b, c
9. Investigation into the effect of a named variable on the rate of respiration of cultures of single-celled organisms	a, b, c, i
10. Investigation into the effect of an environmental variable on the movement of an animal using either a choice chamber or a maze	h
11. Production of a dilution series of a glucose solution and use of colorimetric techniques to produce a calibration curve with which to identify the concentration of glucose in an unknown 'urine' sample	b, c, f
12. Investigation into the effect of a named environmental factor on the distribution of a given species	a, b, h, k, l

What will I learn?

You will learn about a variety of areas some details of which are below and full details in your course outline and specification tick sheet for each topic.

AS and A Level course

Topic 1 – *Biological Molecules All life on Earth shares a common chemistry.*

This provides indirect evidence for evolution. Despite their great variety, the cells of all living organisms contain only a few groups of carbon-based compounds that interact in similar ways.

Carbohydrates are commonly used by cells as respiratory substrates. They also form structural components in plasma membranes and cell walls. Lipids have many uses, including the bilayer of plasma membranes, certain hormones, and as respiratory substrates. Proteins form many cell structures. They are also important as enzymes, chemical messengers, and components of the blood. Nucleic acids carry the genetic code to produce proteins. The genetic code is common to viruses and to all living organisms, providing evidence for evolution.

The most common component of cells is water; hence our search for life elsewhere in the universe involves a search for liquid water.

Topic 2 – *Cells All life on Earth exists as cells.*

These have basic features in common. Differences between cells are due to the addition of extra features. This provides indirect evidence for evolution. All cells arise from other cells, by binary fission in prokaryotic cells and by mitosis and meiosis in eukaryotic cells.

All cells have a cell-surface membrane, the plasma membrane, and, in addition, eukaryotic cells have internal membranes. The basic structure of these membranes is the same and enables control of the passage of substances across exchange surfaces by passive or active transport. Cell-surface membranes contain embedded proteins. Some of these are involved in cell signalling – communication between cells. Others act as antigens, allowing recognition of 'self' and 'foreign' cells by the immune system. Interactions between different types of cells are involved in disease, recovery from disease and prevention of symptoms occurring later if exposed to the same antigen, or antigen-bearing pathogen.

Topic 3 – *Organism Exchange The internal environment of a cell or organism is different from its external environment.*

The exchange of substances between the internal and external environments takes place at exchange surfaces. To truly enter or leave an organism, most substances must cross cell plasma membranes.

In large multicellular organisms, the immediate environment of cells is some form of tissue fluid. Most cells are too far away from exchange surfaces, and from each other, for simple diffusion alone to maintain the composition of tissue fluid within a suitable metabolic range. In large organisms, exchange surfaces are associated with mass transport systems that carry substances between the exchange surfaces and the rest of the body and between parts of the body.

Mass transport maintains the final diffusion gradients that bring substances to and from the cell membranes of individual cells. It also helps to maintain the relatively stable environment that is tissue fluid.

Topic 4 – Genetic information and relationships between organisms

Biological diversity – biodiversity – is reflected in the vast number of species of organisms, in the variation of individual characteristics within a single species and in the variation of cell types within a single multicellular organism.

Differences between species reflect genetic differences. Differences between individuals within a species could be the result of genetic factors, of environmental factors, or a combination of both. A gene is a section of DNA located at a particular site on a DNA molecule, called its locus. The base sequence of each gene carries the genetic code that determines the sequence of amino acids during protein synthesis. The genetic code is the same in all organisms, providing indirect evidence for evolution. Genetic diversity within a species can be caused by gene mutation, chromosome mutation or random factors associated with meiosis and fertilisation. This genetic diversity is acted upon by natural selection, resulting in species becoming better adapted to their environment.

Variation within a species can be measured using differences in the base sequence of DNA or in the amino acid sequence of proteins. Biodiversity within a community can be measured using species richness and an index of diversity.

Topic 5 – Energy transfers in and between organisms Life depends on continuous transfers of energy.

In photosynthesis, light is absorbed by chlorophyll, and this is linked to the production of ATP. In respiration, various substances are used as respiratory substrates. The hydrolysis of these respiratory substrates is linked to the production of ATP. In both respiration and photosynthesis, ATP production occurs when protons diffuse down an electrochemical gradient through molecules of the enzyme ATP synthase, embedded in the membranes of cellular organelles. The process of photosynthesis is common in all photoautotrophic organisms and the process of respiration is common in all organisms, providing indirect evidence for evolution.

In communities, the biological molecules produced by photosynthesis are consumed by other organisms, including animals, bacteria, and fungi. Some of these are used as respiratory substrates by these consumers. Photosynthesis and respiration are not 100% efficient. The transfer of biomass and its stored chemical energy in a community from one organism to a consumer is also not 100% efficient.

Topic 6 – Organisms respond to changes in their internal and external environments.

A stimulus is a change in the internal or external environment. A receptor detects a stimulus. A coordinator formulates a suitable response to a stimulus. An effector produces a response. Receptors are specific to one type of stimulus. Nerve cells pass electrical impulses along their length. A nerve impulse is specific to a target cell only because it releases a chemical messenger directly onto it, producing a response that is usually rapid, short-lived, and localised.

In contrast, mammalian hormones stimulate their target cells via the blood system. They are specific to the tertiary structure of receptors on their target cells and produce responses that are usually slow, long-lasting, and widespread. Plants control their response using hormone-like growth substances.

Topic 7 – Genetics, populations, evolution, and ecosystems the theory of evolution underpins modern Biology.

All new species arise from an existing species. This results in different species sharing a common ancestry, as represented in phylogenetic classification. Common ancestry can explain the similarities between all living organisms, such as common chemistry (eg all proteins made from the same 20 or so amino acids), physiological pathways (eg anaerobic respiration), cell structure, DNA as the genetic

material and a 'universal' genetic code. The individuals of a species share the same genes but (usually) different combinations of alleles of these genes. An individual inherits alleles from their parent or parents.

A species exists as one or more populations. There is variation in the phenotypes of organisms in a population, due to genetic and environmental factors. Two forces affect genetic variation in populations: genetic drift and natural selection. Genetic drift can cause changes in allele frequency in small populations. Natural selection occurs when alleles that enhance the fitness of the individuals that carry them rise in frequency. A change in the allele frequency of a population is evolution.

If a population becomes isolated from other populations of the same species, there will be no gene flow between the isolated population and the others. This may lead to the accumulation of genetic differences in the isolated population, compared with the other populations. These differences may ultimately lead to organisms in the isolated population becoming unable to breed and produce fertile offspring with organisms from the other populations. This reproductive isolation means that a new species has evolved. Populations of different species live in communities. Competition occurs within and between these populations for the means of survival. Within a single community, one population is affected by other populations, the biotic factors, in its environment. Populations within communities are also affected by, and in turn affect, the abiotic (physicochemical) factors in an ecosystem.


Topic 8 – *The control of gene expression.*

Cells can control their metabolic activities by regulating the transcription and translation of their genome. Although the cells within an organism carry the same genetic code, they translate only part of it. In multicellular organisms, this control of translation enables cells to have specialised functions, forming tissues and organs. There are many factors that control the expression of genes and, thus, the phenotype of organisms. Some are external, environmental factors, others are internal factors. The expression of genes is not as simple as once thought, with epigenetic regulation of transcription being increasingly recognised as important.

Humans are learning how to control the expression of genes by altering the epigenome, and how to alter genomes and proteomes of organisms. This has many medical and technological applications. Consideration of cellular control mechanisms underpins the content of this section. Students who have studied it should develop an understanding of the ways in which organisms and cells control their activities. This should lead to an appreciation of common ailments resulting from a breakdown of these control mechanisms and the use of DNA technology in the diagnosis and treatment of human diseases.

Biology A level Specification

Biological molecules

MONOMERS & POLYMERS			
The biochemical basis of life is similar for all living things			
Monomers are the smaller units from which larger molecules are made			
Polymers are molecules made from a large number of monomers joined together			
Monosaccharides, amino acids and nucleotides are examples of monomers			
A condensation reaction joins two molecules together with the formation of a chemical bond and involves the elimination of a molecule of water			
A hydrolysis reaction breaks a chemical bond between two molecules and involves the use of a water molecule			
Monosaccharides are the monomers from which larger carbohydrates are made. Glucose, galactose and fructose are common monosaccharides			
A condensation reaction between two monosaccharides forms a glycosidic bond.			
Disaccharides are formed by the condensation of two monosaccharides			
• maltose is a disaccharide formed by condensation of two glucose molecules			
• sucrose is a disaccharide formed by condensation of a glucose molecule and a fructose molecule			
• lactose is a disaccharide formed by condensation of a glucose molecule and a galactose molecule.			
Glucose has two isomers, α-glucose and β-glucose, with structures:  α-glucose β-glucose			
Polysaccharides are formed by the condensation of many glucose units			
• glycogen and starch are formed by the condensation of α-glucose			
• cellulose is formed by the condensation of β-glucose			
The basic structure and functions of glycogen, starch and cellulose. The relationship of structure to function of these substances in animal cells and plant cells			
Biochemical tests using Benedict's solution for reducing sugars and non-reducing sugars and iodine/potassium iodide for starch			


LIPIDS			
Triglycerides and phospholipids are two groups of lipid			
Triglycerides are formed by the condensation of one molecule of glycerol and three molecules of fatty acid			
A condensation reaction between glycerol and a fatty acid (RCOOH) forms an ester bond			
The R-group of a fatty acid may be saturated or unsaturated			
In phospholipids, one of the fatty acids of a triglyceride is substituted by a phosphate-containing group			
The different properties of triglycerides and phospholipids related to their different structures			
The emulsion test for lipids			
Students should be able to recognise, from diagrams, saturated & unsaturated fatty acids			
Students should be able to explain the different properties of triglycerides and phospholipids			

PROTEINS			
General properties of proteins			
Amino acids are the monomers from which proteins are made. The general structure of an amino acid as: $ \begin{array}{c} \text{R} \\ \\ \text{H}_2\text{N} - \text{C} - \text{COOH} \\ \\ \text{H} \end{array} $ where NH₂ represents an amine group, COOH represents a carboxyl group and R represents a side chain. The twenty amino acids that are common in all organisms differ only in their side group			
A condensation reaction between two amino acids forms a peptide bond			
• dipeptides are formed by the condensation of two amino acids			
• polypeptides are formed by the condensation of many amino acids			
A functional protein may contain one or more polypeptides			
The role of hydrogen bonds, ionic bonds and disulfide bridges in the structure of proteins			

Proteins have a variety of functions within all living organisms. The relationship between primary, secondary, tertiary and quaternary structure, and protein function			
Biuret test for proteins			
Students should be able to relate the structure of proteins to properties of proteins named throughout the specification			
Many proteins are enzymes			
Each enzyme lowers the activation energy of the reaction it catalyses			
The induced-fit model of enzyme action.			
The properties of an enzyme relate to the tertiary structure of its active site and its ability to combine with complementary substrate(s) to form an enzyme-substrate complex			
The specificity of enzymes			
The effects of the following factors on the rate of enzyme controlled reactions – enzyme concentration, substrate concentration, concentration of competitive and of noncompetitive inhibitors, pH and temperature			
Students should be able to appreciate how models of enzyme action have changed over time			
Students should be able to appreciate that enzymes catalyse a wide range of intracellular and extracellular reactions that determine structures and functions from cellular to whole-organism level			
<i>Required practical 1: Investigation into the effect of a named variable on the rate of an enzyme-controlled reaction.</i>			
<i>Students could identify the variables that must be controlled in their investigation into rate of reaction</i>			
<i>Students could calculate the uncertainty of their measurements of the rate of reaction</i>			
<i>Students could select an appropriate format for the graphical presentation of the results of their investigation into the rate of enzyme controlled reactions</i>			
<i>Students could use a tangent to find the initial rate of an enzyme-controlled reaction</i>			

NUCLEIC ACIDS ARE IMPORTANT INFORMATION-CARRYING MOLECULES			
Structure of DNA and RNA			
Deoxyribonucleic acid (DNA) and ribonucleic acid (RNA) are important information-carrying molecules. In all living cells, DNA holds genetic information and RNA transfers genetic information from DNA to the ribosomes			
Ribosomes are formed from RNA and proteins			

Both DNA and RNA are polymers of nucleotides. Each nucleotide is formed from a pentose, a nitrogen-containing organic base and a phosphate group:



- The components of a DNA nucleotide are deoxyribose, a phosphate group and one of the organic bases adenine, cytosine, guanine or thymine.
- The components of an RNA nucleotide are ribose, a phosphate group and one of the organic bases adenine, cytosine, guanine or uracil.
- A condensation reaction between two nucleotides forms a phosphodiester bond.

A DNA molecule is a double helix with two polynucleotide chains held together by hydrogen bonds between specific complementary base pairs

An RNA molecule is a relatively short polynucleotide chain

Students should be able to appreciate that the relative simplicity of DNA led many scientists to doubt that it carried the genetic code


DNA replication

The semi-conservative replication of DNA ensures genetic continuity between generations of cells

The process of semi-conservative replication of DNA in terms of:

- unwinding of the double helix
- breakage of hydrogen bonds between complementary bases in the polynucleotide strands
- the role of DNA helicase in unwinding DNA and breaking its hydrogen bonds
- attraction of new DNA nucleotides to exposed bases on template strands and base pairing
- the role of DNA polymerase in the condensation reaction that joins adjacent nucleotides

Students should be able to evaluate the work of scientists in validating the Watson–Crick model of DNA replication

ATP			
A single molecule of adenosine triphosphate (ATP) is a nucleotide derivative and is formed from a molecule of ribose, a molecule of adenine and three phosphate groups 			
Hydrolysis of ATP to adenosine diphosphate (ADP) and an inorganic phosphate group (P_i) is catalysed by the enzyme ATP hydrolase.			
The hydrolysis of ATP can be coupled to energy-requiring reactions within cells.			
The inorganic phosphate released during the hydrolysis of ATP can be used to phosphorylate other compounds, often making them more reactive.			
ATP is resynthesised by the condensation of ADP and P_i . This reaction is catalysed by the enzyme ATP synthase during photosynthesis, or during respiration			
Water is a major component of cells. It has several properties that are important in biology. In particular, water:			
is a metabolite in many metabolic reactions, including condensation and hydrolysis reactions			
- is an important solvent in which metabolic reactions occur - has a relatively high heat capacity, buffering changes in temperature			
has a relatively large latent heat of vaporisation, providing a cooling effect with little loss of water through evaporation			
has strong cohesion between water molecules; this supports columns of water in the tube-like transport cells of plants and produces surface tension where water meets air			

INORGANIC IONS			
Inorganic ions occur in solution in the cytoplasm and body fluids of organisms, some in high concentrations and others in very low concentrations			
Each type of ion has a specific role, depending on its properties			
Students should be able to recognise the role of ions in the following topics: hydrogen ions and pH; iron ions as a component of haemoglobin; sodium ions in the co-transport of glucose and amino acids; and phosphate ions as components of DNA and of ATP			

Cells

CELL STRUCTURE			
Structure of eukaryotic cells			
The structure of eukaryotic cells, restricted to the structure and function of:			
• cell-surface membrane			
• nucleus (containing chromosomes, consisting of protein-bound, linear DNA, and one or more nucleoli)			
• mitochondria			
• chloroplasts (in plants and algae)			
• golgi apparatus and golgi vesicles			
• ribosomes			
• rough endoplasmic reticulum and smooth endoplasmic reticulum			
• cell wall (in plants, algae and fungi)			
• lysosomes (a membrane-bound organelle that releases hydrolytic enzymes)			
• cell vacuole (in plants)			
In complex multicellular organisms, eukaryotic cells become specialised for specific functions. Specialised cells are organised into tissues, tissues into organs and organs into systems			
Be able to apply their knowledge of these features in explaining adaptations of eukaryotic cells			
Structure of prokaryotic cells and of viruses			
Prokaryotic cells are much smaller than eukaryotic cells. They also differ from eukaryotic cells in having:			
• cytoplasm that lacks membrane-bound organelles			
• smaller ribosomes			
• no nucleus; instead they have a single circular DNA molecule that is free in the cytoplasm and is not associated with proteins			
• a cell wall that contains murein, a glycoprotein			
In addition, many prokaryotic cells have:			
• one or more plasmids			
• a capsule surrounding the cell			
• one or more flagella.			

Details of these structural differences are not required			
Viruses are acellular and non-living. The structure of virus particles to include genetic material, capsid and attachment protein			
Methods of studying cells			
The principles and limitations of optical microscopes, transmission electron microscopes and scanning electron microscopes			
Measuring the size of an object viewed with an optical microscope. The difference between magnification and resolution			
Use of the formula: magnification = size of image / size of real object			
Principles of cell fractionation and ultracentrifugation as used to separate cell components			
Be able to appreciate that there was a considerable period of time during which the scientific community distinguished between artefacts and cell organelles			

ALL CELLS ARISE FROM OTHER CELLS			
Within multicellular organisms, not all cells retain the ability to divide			
Eukaryotic cells that do retain the ability to divide show a cell cycle.			
DNA replication occurs during the interphase of the cell cycle.			
Mitosis is the part of the cell cycle in which a eukaryotic cell divides to produce two daughter cells, each with the identical copies of DNA produced by the parent cell during DNA replication			
The behaviour of chromosomes during interphase, prophase, metaphase, anaphase and telophase of mitosis. The role of spindle fibres attached to centromeres in the separation of chromatids			
Division of the cytoplasm (cytokinesis) usually occurs, producing two new cells			
Be able to recognise the stages of the cell cycle: interphase, prophase, metaphase, anaphase and telophase (including cytokinesis)			
Be able to explain the appearance of cells in each stage of mitosis			
Mitosis is a controlled process. Uncontrolled cell division can lead to the formation of tumours and of cancers. Many cancer treatments are directed at controlling the rate of cell division			
Binary fission in prokaryotic cells involves:			
replication of the circular DNA and of plasmids			
division of the cytoplasm to produce two daughter cells, each with a single copy of the circular DNA and a variable number of copies of plasmids			
Being non-living, viruses do not undergo cell division. Following injection of their nucleic acid, the infected host cell replicates the virus particles			

<i>Required practical 2: Preparation of stained squashes of cells from plant root tips; set-up and use of an optical microscope to identify the stages of mitosis in these stained squashes and calculation of a mitotic index. Students should measure the apparent size of cells in the root tip and calculate their actual size using the formula</i>			
--	--	--	--

TRANSPORT ACROSS CELL MEMBRANES			
The basic structure of all cell membranes, including cell-surface membranes and the membranes around the cell organelles of eukaryotes, is the same			
The arrangement and any movement of phospholipids, proteins, glycoproteins and glycolipids in the fluid-mosaic model of membrane structure. Cholesterol may also be present in cell membranes where it restricts the movement of other molecules making up the membrane.			
Movement across membranes occurs by:			
• simple diffusion (involving limitations imposed by the nature of the phospholipid bilayer)			
• facilitated diffusion (involving the roles of carrier proteins and channel proteins)			
• osmosis (explained in terms of water potential)			
• active transport (involving the role of carrier proteins and the importance of the hydrolysis of ATP)			
• co-transport (illustrated by the absorption of sodium ions and glucose by cells lining the mammalian ileum)			
Cells may be adapted for rapid transport across their internal or external membranes by an increase in surface area of, or by an increase in the number of protein channels and carrier molecules in, their membranes			
Students should be able to: • explain the adaptations of specialised cells in relation to the rate of transport across their internal and external membranes • explain how surface area, number of channel or carrier proteins and differences in gradients of concentration or water potential affect the rate of movement across cell membranes.			
<i>Required practical 3: Production of a dilution series of a solute to produce a calibration curve with which to identify the water potential of plant tissue</i>			
<i>Required practical 4: Investigation into the effect of a named variable on the permeability of cell-surface membranes</i>			

CELL RECOGNITION AND THE IMMUNE SYSTEM			
Each type of cell has specific molecules on its surface that identify it. These molecules include proteins and enable the immune system to identify:			
• pathogens			
• cells from other organisms of the same species			

• abnormal body cells			
• toxins			
Definition of antigen. The effect of antigen variability on disease and disease prevention			
Phagocytosis of pathogens. The subsequent destruction of ingested pathogens by lysozymes			
The response of T lymphocytes to a foreign antigen (the cellular response). • The role of antigen-presenting cells in the cellular response. • The role of helper T cells (TH cells) in stimulating cytotoxic T cells (TC cells), B cells and phagocytes. The role of other T cells is not required.			
The response of B lymphocytes to a foreign antigen, clonal selection and the release of monoclonal antibodies (the humoral response).			
• Definition of antibody.			
• Antibody structure.			
• The formation of an antigen-antibody complex, leading to the destruction of the antigen, limited to agglutination and phagocytosis of bacterial cells.			
• The roles of plasma cells and of memory cells in producing primary and secondary immune responses			
The use of vaccines to provide protection for individuals and populations against disease. The concept of herd immunity			
The differences between active and passive immunity			
Structure of the human immunodeficiency virus (HIV) and its replication in helper T cells			
How HIV causes the symptoms of AIDS. Why antibiotics are ineffective against viruses			
The use of monoclonal antibodies in:			
• medical diagnosis			
• targeting medication to specific cell types by attaching a therapeutic drug to an antibody			
• medical diagnosis			
Details of the commercial or scientific production of monoclonal antibodies are not required			
Ethical issues associated with the use of vaccines and monoclonal antibodies			
The use of antibodies in the ELISA test			
Students should be able to:			
• discuss ethical issues associated with the use of vaccines and monoclonal antibodies			
• evaluate methodology, evidence and data relating to the use of vaccines and monoclonal antibodies			

Organisms exchange substances with their environment

SURFACE AREA TO VOLUME RATIO			
The relationship between the size of an organism or structure and its surface area to volume ratio			
Changes to body shape and the development of systems in larger organisms as adaptations that facilitate exchange as this ratio reduces			
Students should be able to appreciate the relationship between surface area to volume ratio and metabolic rate			

GAS EXCHANGE			
Adaptations of gas exchange surfaces, shown by gas exchange:			
across the body surface of a single-celled organism			
in the tracheal system of an insect (tracheae, tracheoles and spiracles)			
across the gills of fish (gill lamellae and filaments including the counter-current principle)			
by the leaves of dicotyledonous plants (mesophyll and stomata)			
Structural and functional compromises between the opposing needs for efficient gas exchange and the limitation of water loss shown by terrestrial insects and xerophytic plants			
The gross structure of the human gas exchange system limited to the alveoli, bronchioles, bronchi, trachea and lungs			
The essential features of the alveolar epithelium as a surface over which gas exchange takes place			
Ventilation and the exchange of gases in the lungs. The mechanism of breathing to include the role of the diaphragm and the antagonistic interaction between the external and internal intercostal muscles in bringing about pressure changes in the thoracic cavity			
Students should be able to:			
interpret information relating to the effects of lung disease on gas exchange and/or ventilation			
interpret data relating to the effects of pollution and smoking on the incidence of lung disease			
analyse and interpret data associated with specific risk factors and the incidence of lung disease			
evaluate the way in which experimental data led to statutory restrictions on the sources of risk factors			
recognise correlations and causal relationships			

DIGESTION AND ABSORPTION			
During digestion, large biological molecules are hydrolysed to smaller molecules that can be absorbed across cell membranes			
Digestion in mammals of:			
• carbohydrates by amylases and membrane-bound disaccharidases			
• lipids by lipase, including the action of bile salts			
• proteins by endopeptidases, exopeptidases and membrane bound dipeptidases.			
Mechanisms for the absorption of the products of digestion by cells lining the ileum of mammals, to include:			
• co-transport mechanisms for the absorption of amino acids and of monosaccharides			
• the role of micelles in the absorption of lipids			

Mass transport

MASS TRANSPORT			
Mass transport in animals			
The haemoglobins are a group of chemically similar molecules found in many different organisms. Haemoglobin is a protein with a quaternary structure			
The role of haemoglobin and red blood cells in the transport of oxygen. The loading, transport and unloading of oxygen in relation to the oxyhaemoglobin dissociation curve. The cooperative nature of oxygen binding to show that the change in shape of haemoglobin caused by binding of the first oxygens makes the binding of further oxygens easier. The effects of carbon dioxide concentration on the dissociation of oxyhaemoglobin (the Bohr effect)			
Many animals are adapted to their environment by possessing different types of haemoglobin with different oxygen transport properties			
The general pattern of blood circulation in a mammal. Names are required only of the coronary arteries and of the blood vessels entering and leaving the heart, lungs and kidneys			
The gross structure of the human heart. Pressure and volume changes and associated valve movements during the cardiac cycle that maintain a unidirectional flow of blood			
The structure of arteries, arterioles and veins in relation to their function			
The structure of capillaries and the importance of capillary beds as exchange surfaces. The formation of tissue fluid and its return to the circulatory system			
Students should be able to:			
• analyse and interpret data relating to pressure and volume changes during the cardiac cycle			
• analyse and interpret data associated with specific risk factors and the incidence of			

cardiovascular disease			
• evaluate conflicting evidence associated with risk factors affecting cardiovascular disease			
• recognise correlations and causal relationships			
<i>Required practical 5: Dissection of animal or plant gas exchange system or mass transport system or of organ within such a system</i>			
Mass transport in plants			
Xylem: the tissue that transports water in the stem and leaves of plants. The cohesion-tension theory of water transport in the xylem			
Phloem: the tissue that transports organic substances in plants. The mass flow hypothesis for the mechanism of translocation in plants. The use of tracers and ringing experiments to investigate transport in plants			
Students should be able to:			
• recognise correlations and causal relationships			
• interpret evidence from tracer and ringing experiments and to evaluate the evidence for and against the mass flow hypothesis			

Genetic Information, Variation and Relationships Between Organisms

DNA, GENES AND CHROMOSOMES			
In prokaryotic cells, DNA molecules are short, circular and not associated with proteins			
In the nucleus of eukaryotic cells, DNA molecules are very long, linear and associated with proteins, called histones. Together a DNA molecule and its associated proteins form a chromosome			
The mitochondria and chloroplasts of eukaryotic cells also contain DNA which, like the DNA of prokaryotes, is short, circular and not associated with protein			
A gene is a base sequence of DNA that codes for:			
• the amino acid sequence of a polypeptide			
• a functional RNA (including ribosomal RNA and tRNAs)			
A gene occupies a fixed position, called a locus, on a particular DNA molecule			
A sequence of three DNA bases, called a triplet, codes for a specific amino acid. The genetic code is universal, non-overlapping and degenerate			
In eukaryotes, much of the nuclear DNA does not code for polypeptides. There are, for example, non-coding multiple repeats of base sequences between genes. Even within a gene only some sequences, called exons, code for amino acid sequences. Within the gene, these exons are separated by one or more non-coding sequences, called introns			

DNA AND PROTEIN SYNTHESIS			
The concept of the genome as the complete set of genes in a cell and of the proteome as the full range of proteins that a cell is able to produce			
The structure of molecules of messenger RNA (mRNA) and of transfer RNA (tRNA)			
Transcription as the production of mRNA from DNA. The role of RNA polymerase in joining mRNA nucleotides			
• In prokaryotes, transcription results directly in the production of mRNA from DNA.			
• In eukaryotes, transcription results in the production of premRNA; this is then spliced to form mRNA			
Translation as the production of polypeptides from the sequence of codons carried by mRNA. The roles of ribosomes, tRNA and ATP			
Be able to relate the base sequence of nucleic acids to the amino acid sequence of polypeptides, when provided with suitable data about the genetic code			
Be able to interpret data from experimental work investigating the role of nucleic acids			
Students will not be required to recall in written papers specific codons and the amino acids for which they code			

GENETIC DIVERSITY CAN ARISE AS A RESULT OF MUTATION OR DURING MEIOSIS			
Gene mutations involve a change in the base sequence of chromosomes. They can arise spontaneously during DNA replication and include base deletion and base substitution. Due to the degenerate nature of the genetic code, not all base substitutions cause a change in the sequence of encoded amino acids. Mutagenic agents can increase the rate of gene mutation			
Mutations in the number of chromosomes can arise spontaneously by chromosome non-disjunction during meiosis			
Meiosis produces daughter cells that are genetically different from each other			
The process of meiosis only in sufficient detail to show how:			
• two nuclear divisions result usually in the formation of four haploid daughter cells from a single diploid parent cell			
• genetically different daughter cells result from the independent segregation of homologous chromosomes			
• crossing over between homologous chromosomes results in further genetic variation among daughter cells			
Be able to:			

• complete diagrams showing the chromosome content of cells after the first and second meiotic division, when given the chromosome content of the parent cell			
• explain the different outcome of mitosis and meiosis			
• recognise where meiosis occurs when given information about an unfamiliar life cycle			
• explain how random fertilisation of haploid gametes further increases genetic variation within a species			

GENETIC DIVERSITY AND ADAPTATION			
Genetic diversity as the number of different alleles of genes in a population			
Genetic diversity is a factor enabling natural selection to occur			
The principles of natural selection in the evolution of populations.			
• Random mutation can result in new alleles of a gene			
• Many mutations are harmful but, in certain environments, the new allele of a gene might benefit its possessor, leading to increased reproductive success			
• The advantageous allele is inherited by members of the next generation			
• As a result, over many generations, the new allele increases in frequency in the population			
Directional selection, exemplified by antibiotic resistance in bacteria, and stabilising selection, exemplified by human birth weights.			
Natural selection results in species that are better adapted to their environment. These adaptations may be anatomical, physiological or behavioural			
Be able to:			
• use unfamiliar information to explain how selection produces changes within a population of a species			
• interpret data relating to the effect of selection in producing change within populations			
• show understanding that adaptation and selection are major factors in evolution and contribute to the diversity of living organisms			
<i>Required practical 6: Use of aseptic techniques to investigate the effect of antimicrobial substances on microbial growth</i>			

SPECIES AND TAXONOMY			
Two organisms belong to the same species if they are able to produce fertile offspring. Courtship behaviour as a necessary precursor to successful mating. The role of courtship in species recognition			
A phylogenetic classification system attempts to arrange species into groups based on their evolutionary origins and relationships. It uses a hierarchy in which smaller groups are placed within larger groups, with no overlap between groups. Each group is called a taxon (plural taxa)			
One hierarchy comprises the taxa: domain, kingdom, phylum, class, order, family, genus and species			
Each species is universally identified by a binomial consisting of the name of its genus and species, eg, Homo sapiens			
Recall of different taxonomic systems, such as the three domain or five kingdom systems, will not be required			
Students should be able to appreciate that advances in immunology and genome sequencing help to clarify evolutionary relationships between organisms			

BIODIVERSITY WITHIN A COMMUNITY			
Biodiversity can relate to a range of habitats, from a small local habitat to the Earth			
Species richness is a measure of the number of different species in a community			
An index of diversity describes the relationship between the number of species in a community and the number of individuals in each species			
Calculation of an index of diversity (d) from the formula $d = \frac{N(N-1)}{\sum n(n-1)}$ Where N = total number of organisms of all species and n = total number of organisms of each species			
Farming techniques reduce biodiversity. The balance between conservation and farming			

INVESTIGATING DIVERSITY			
Genetic diversity within, or between species, can be made by comparing:			
• the frequency of measurable or observable characteristics			
• the base sequence of DNA			

• the base sequence of mRNA			
• the amino acid sequence of the proteins encoded by DNA and mRNA.			
Be able to interpret data relating to similarities and differences in the base sequences of DNA and in the amino acid sequences of proteins to suggest relationships between different organisms within a species and between species			
Be able to appreciate that gene technology has caused a change in the methods of investigating genetic diversity; inferring DNA differences from measurable or observable characteristics has been replaced by direct investigation of DNA sequences			
Knowledge of gene technologies will not be tested			
Quantitative investigations of variation within a species involve:			
• collecting data from random samples			
• calculating a mean value of the collected data and the standard deviation of that mean			
• interpreting mean values and their standard deviations			
Students will not be required to calculate standard deviations in written papers			

Energy Transfers in and Between Organisms (A-level Only)

PHOTOSYNTHESIS (A2)			
The light dependent reaction in such detail as to show that:			
• chlorophyll absorbs light, leading to photoionisation of chlorophyll			
• some of the energy from electrons released during photoionisation is conserved in the production of ATP and reduced NADP			
• the production of ATP involves electron transfer associated with the transfer of electrons down the electron transfer chain and passage of protons across chloroplast membranes and is catalysed by ATP synthase embedded in these membranes (chemiosmotic theory)			
• photolysis of water produces protons, electrons and oxygen			
The light-independent reaction uses reduced NADP from the light-dependent reaction to form a simple sugar. The hydrolysis of ATP, also from the light-dependent reaction, provides the additional energy for this reaction			
The light-independent reaction in such detail as to show that:			
• carbon dioxide reacts with ribulose biphosphate (RuBP) to form two molecules of glycerate 3-phosphate (GP). This reaction is catalysed by the enzyme rubisco			

• ATP and reduced NADP from the light-dependent reaction are used to reduce GP to triose phosphate			
• some of the triose phosphate is used to regenerate RuBP in the Calvin cycle			
• some of the triose phosphate is converted to useful organic substances			
Be able to identify environmental factors that limit the rate of photosynthesis			
Be able to evaluate data relating to common agricultural practices used to overcome the effect of these limiting factors			
<i>Required practical 7: Use of chromatography to investigate the pigments isolated from leaves of different plants, eg, leaves from shade-tolerant and shade-intolerant plants or leaves of different colours</i>			
<i>Required practical 8: Investigation into the effect of a named factor on the rate of dehydrogenase activity in extracts of chloroplasts</i>			

RESPIRATION (A2)			
Respiration produces ATP			
Glycolysis is the first stage of anaerobic and aerobic respiration. It occurs in the cytoplasm and is an anaerobic process			
Glycolysis involves the following stages:			
• phosphorylation of glucose to glucose phosphate, using ATP			
• production of triose phosphate			
• oxidation of triose phosphate to pyruvate with a net gain of ATP and reduced NAD			
If respiration is only anaerobic, pyruvate can be converted to ethanol or lactate using reduced NAD. The oxidised NAD produced in this way can be used in further glycolysis			
If respiration is aerobic, pyruvate from glycolysis enters the mitochondrial matrix by active transport			
Aerobic respiration in such detail as to show that:			
• pyruvate is oxidised to acetate, producing reduced NAD in the process			
• acetate combines with coenzyme A in the link reaction to produce acetylcoenzyme A			
• acetyl coenzyme A reacts with a four-carbon molecule, releasing coenzyme A and producing a six-carbon molecule that enters the Krebs cycle			
• in a series of oxidation-reduction reactions, the Krebs cycle generates reduced coenzymes and ATP by substrate-level phosphorylation, and carbon dioxide is lost			
• synthesis of ATP by oxidative phosphorylation is associated with the transfer of electrons down the electron transfer chain and passage of protons across inner mitochondrial membranes and is catalysed by ATP synthase embedded in these membranes			

(chemiosmotic theory)			
other respiratory substrates include the breakdown products of lipids and amino acids, which enter the Krebs cycle			
<i>Required practical 9: Investigation into the effect of a named variable on the rate of respiration of cultures of single-celled organisms</i>			

ENERGY AND ECOSYSTEMS (A2)			
In any ecosystem, plants synthesise organic compounds from atmospheric, or aquatic, carbon dioxide			
Most of the sugars synthesised by plants are used by the plant as respiratory substrates. The rest are used to make other groups of biological molecules. These biological molecules form the biomass of the plants			
Biomass can be measured in terms of mass of carbon or dry mass of tissue per given area. The chemical energy store in dry biomass can be estimated using calorimetry			
Gross primary production (<i>GPP</i>) is the chemical energy store in plant biomass, in a given area or volume			
Net primary production (<i>NPP</i>) is the chemical energy store in plant biomass after respiratory losses to the environment have been taken into account, ie $NPP = GPP - R$ where <i>GPP</i> represents gross production and <i>R</i> represents respiratory losses to the environment.			
This net primary production is available for plant growth and reproduction. It is also available to other trophic levels in the ecosystem, such as herbivores and decomposers			
The net production of consumers (<i>N</i>), such as animals, can be calculated as: $N = I - (F + R)$ where <i>I</i> represents the chemical energy store in ingested food, <i>F</i> represents the chemical energy lost to the environment in faeces and urine and <i>R</i> represents the respiratory losses to the environment			
Primary and secondary productivity is the rate of primary or secondary production, respectively. It is measured as biomass in a given area in a given time eg $\text{kJ ha}^{-1} \text{ year}^{-1}$			
Be able to appreciate the ways in which production is affected by farming practices designed to increase the efficiency of energy transfer by:			
simplifying food webs to reduce energy losses to non-human food chains			

• reducing respiratory losses within a human food chain			
---	--	--	--

NUTRIENT CYCLES (A2)			
Nutrients are recycled within natural ecosystems, exemplified by the nitrogen cycle and the phosphorus cycle			
Microorganisms play a vital role in recycling chemical elements such as phosphorus and nitrogen			
• The role of saprobionts in decomposition			
• The role of mycorrhizae in facilitating the uptake of water and inorganic ions by plants			
• The role of bacteria in the nitrogen cycle in sufficient detail to illustrate the processes of saprobiotic nutrition, ammonification, nitrification, nitrogen fixation and denitrification - Names of individual species of bacteria are not required			
The use of natural and artificial fertilisers to replace the nitrates and phosphates lost by harvesting plants and removing livestock			
The environmental issues arising from the use of fertilisers including leaching and eutrophication			

Organisms Respond to Changes in their Internal and External

STIMULI, BOTH INTERNAL AND EXTERNAL, ARE DETECTED AND LEAD TO A RESPONSE			
Survival and response			
Organisms increase their chance of survival by responding to changes in their environment			
In flowering plants, specific growth factors move from growing regions to other tissues, where they regulate growth in response to directional stimuli			
The effect of different concentrations of indoleacetic acid (IAA) on cell elongation in the roots and shoots of flowering plants as an explanation of gravitropism and phototropism in flowering plants			
Taxes and kinesis as simple responses that can maintain a mobile organism in a favourable environment			
The protective effect of a simple reflex, exemplified by a three neurone simple reflex. Details of spinal cord and dorsal and ventral roots are not required			
<i>Required practical 10: Investigation into the effect of an environmental variable on the movement of an animal using either a choice chamber or a maze</i>			
Receptors			
The Pacinian corpuscle should be used as an example of a receptor to illustrate that:			
• receptors respond only to specific stimuli			
• stimulation of a receptor leads to the establishment of a generator potential			

The basic structure of a Pacinian corpuscle			
Deformation of stretch-mediated sodium ion channels in a Pacinian corpuscle leads to the establishment of a generator potential			
The human retina in sufficient detail to show how differences in sensitivity to light, sensitivity to colour and visual acuity are explained by differences in the optical pigments of rods and cones and the connections rods and cones make in the optic nerve			
Control of heart rate			
Myogenic stimulation of the heart and transmission of a subsequent wave of electrical activity. The roles of the sinoatrial node (SAN), atrioventricular node (AVN) and Purkyne tissue in the bundle of His			
The roles and locations of chemoreceptors and pressure receptors and the roles of the autonomic nervous system and effectors in controlling heart rate			

NERVOUS COORDINATION			
Nerve impulses			
The structure of a myelinated motor neurone			
The establishment of a resting potential in terms of differential membrane permeability, electrochemical gradients and the movement of sodium ions and potassium ions			
Changes in membrane permeability lead to depolarisation and the generation of an action potential. The all-or-nothing principle			
The passage of an action potential along non-myelinated and myelinated axons, resulting in nerve impulses			
The nature and importance of the refractory period in producing discrete impulses and in limiting the frequency of impulse transmission			
Factors affecting the speed of conductance: myelination and saltatory conduction; axon diameter; temperature			
Synaptic transmission			
The detailed structure of a synapse and of a neuromuscular junction			
The sequence of events involved in transmission across a cholinergic synapse in sufficient detail to explain:			
• unidirectionality			
• temporal and spatial summation			
• inhibition by inhibitory synapses			
A comparison of transmission across a cholinergic synapse and across a neuromuscular junction			
Be able to use information provided to predict and explain the effects of specific drugs on a synapse			

Recall of the names and mode of action of individual drugs will not be required			
---	--	--	--

SKELETAL MUSCLES ARE STIMULATED TO CONTRACT BY NERVES AND ACT AS EFFECTORS			
Muscles act in antagonistic pairs against an incompressible skeleton			
Gross and microscopic structure of skeletal muscle. The ultrastructure of a myofibril			
The roles of actin, myosin, calcium ions and ATP in myofibril contraction			
The roles of calcium ions and tropomyosin in the cycle of actinomyosin bridge formation			
The role of troponin is not required			
The roles of ATP and phosphocreatine in muscle contraction			
The structure, location and general properties of slow and fast skeletal muscle fibres			

HOMEOSTASIS IS THE MAINTENANCE OF A STABLE INTERNAL ENVIRONMENT			
Principles of homeostasis and negative feedback			
Homeostasis in mammals involves physiological control systems that maintain the internal environment within restricted limits			
The importance of maintaining a stable core temperature and stable blood pH in relation to enzyme activity			
The importance of maintaining a stable blood glucose concentration in terms of availability of respiratory substrate and of the water potential of blood			
Negative feedback restores systems to their original level			
The possession of separate mechanisms involving negative feedback controls departures in different directions from the original state, giving a greater degree of control			
Be able to interpret information relating to examples of negative and positive feedback			
Control of blood glucose concentration			
The factors that influence blood glucose concentration			
The role of the liver in glycogenesis, glycogenolysis and gluconeogenesis			
The action of insulin by:			
• attaching to receptors on the surfaces of target cells			
• controlling the uptake of glucose by regulating the inclusion of channel proteins in the surface membranes of target cells			

• activating enzymes involved in the conversion of glucose to glycogen			
The action of glucagon by:			
• attaching to receptors on the surfaces of target cells			
• activating enzymes involved in the conversion of glycogen to glucose			
• activating enzymes involved in the conversion of glycerol and amino acids into glucose			
The role of adrenaline by:			
• attaching to receptors on the surfaces of target cells			
• activating enzymes involved in the conversion of glycogen to glucose			
The second messenger model of adrenaline and glucagon action, involving adenylate cyclase, cyclic AMP (cAMP) and protein kinase			
The causes of types I and II diabetes and their control by insulin and/or manipulation of the diet			
Be able to evaluate the positions of health advisers and the food industry in relation to the increased incidence of type II diabetes			
<i>Required practical 11: Production of a dilution series of a glucose solution and use of colorimetric techniques to produce a calibration curve with which to identify the concentration of glucose in an unknown 'urine' sample</i>			
Control of blood water potential			
Osmoregulation as control of the water potential of the blood			
The roles of the hypothalamus, posterior pituitary and antidiuretic hormone (ADH) in osmoregulation			
The structure of the nephron and its role in:			
• the formation of glomerular filtrate			
• reabsorption of glucose and water by the proximal convoluted tubule			
• maintaining a gradient of sodium ions in the medulla by the loop of Henle			
• reabsorption of water by the distal convoluted tubule and collecting ducts			

Genetics, Populations, Evolution and Ecosystems (A-level Only)

INHERITANCE			
The genotype is the genetic constitution of an organism			
The phenotype is the expression of this genetic constitution and its interaction with the environment			
There may be many alleles of a single gene			

Alleles may be dominant, recessive or codominant			
In a diploid organism, the alleles at a specific locus may be either homozygous or heterozygous			
The use of fully labelled genetic diagrams to interpret, or predict, the results of:			
• monohybrid and dihybrid crosses involving dominant, recessive and codominant alleles			
• crosses involving sex-linkage, autosomal linkage, multiple alleles and epistasis			
Use of the chi-squared (χ^2) test to compare the goodness of fit of observed phenotypic ratios with expected ratios			

POPULATIONS			
Species exist as one or more populations			
A population as a group of organisms of the same species occupying a particular space at a particular time that can potentially interbreed			
The concepts of gene pool and allele frequency			
The Hardy–Weinberg principle provides a mathematical model, which predicts that allele frequencies will not change from generation to generation. The conditions under which the principle applies			
The frequency of alleles, genotypes and phenotypes in a population can be calculated using the Hardy–Weinberg equation: $p^2 + 2pq + q^2 = 1$ where p is the frequency of one (usually the dominant) allele and q is the frequency of the other (usually recessive) allele of the gene			

EVOLUTION MAY LEAD TO SPECIATION			
Individuals within a population of a species may show a wide range of variation in phenotype. This is due to genetic and environmental factors. The primary source of genetic variation is mutation. Meiosis and the random fertilisation of gametes during sexual reproduction produce further genetic variation			
Predation, disease and competition for the means of survival result in differential survival and reproduction, ie natural selection			
Those organisms with phenotypes providing selective advantages are likely to produce more offspring and pass on their favourable alleles to the next generation. The effect of this differential reproductive success on the allele frequencies within a gene pool			
The effects of stabilising, directional and disruptive selection			

Evolution as a change in the allele frequencies in a population			
Reproductive separation of two populations can result in the accumulation of difference in their gene pools. New species arise when these genetic differences lead to an inability of members of the populations to interbreed and produce fertile offspring. In this way, new species arise from existing species			
Allopatric and sympatric speciation			
The importance of genetic drift in causing changes in allele frequency in small populations			
Be able to explain why individuals within a population of a species may show a wide range of variation in phenotype			
Be able to explain why genetic drift is important only in small populations			
Be able to explain how natural selection and isolation may result in change in the allele and phenotype frequency and lead to the formation of a new species			
Be able to explain how evolutionary change over a long period of time has resulted in a great diversity of species			

POPULATIONS IN ECOSYSTEMS			
Populations of different species form a community. A community and the non-living components of its environment together form an ecosystem. Ecosystems can range in size from the very small to the very large			
Within a habitat, a species occupies a niche governed by adaptation to both abiotic and biotic conditions			
An ecosystem supports a certain size of population of a species, called the carrying capacity. This population size can vary as a result of:			
• the effect of abiotic factors			
• interactions between organisms: interspecific and intraspecific competition and predation			
The size of a population can be estimated using:			
• randomly placed quadrats, or quadrats along a belt transect, for slow-moving or non-motile organisms			
• the mark-release-recapture method for motile organisms. The assumptions made when using the mark-release-recapture method			
Ecosystems are dynamic systems			
Primary succession, from colonisation by pioneer species to climax community			
At each stage in succession, certain species may be recognised which change the environment so that it becomes more suitable for other species with different adaptations. The new species may change the environment in such a way that it becomes less suitable for the previous species			

Changes that organisms produce in their abiotic environment can result in a less hostile environment and change biodiversity			
Conservation of habitats frequently involves management of succession			
Be able to show understanding of the need to manage the conflict between human needs and conservation in order to maintain the sustainability of natural resources			
Be able to evaluate evidence and data concerning issues relating to the conservation of species and habitats and consider conflicting evidence			
Be able to use given data to calculate the size of a population estimated using the mark-release-recapture method			
<i>Required practical 12: Investigation into the effect of a named environmental factor on the distribution of a given species</i>			

The Control of Gene Expression (A-level Only)

ALTERATION OF THE SEQUENCE OF BASES IN DNA CAN ALTER THE STRUCTURE OF PROTEINS			
Gene mutations might arise during DNA replication. They include addition, deletion, substitution, inversion, duplication and translocation of bases			
Gene mutations occur spontaneously. The mutation rate is increased by mutagenic agents. Mutations can result in a different amino acid sequence in the encoded polypeptide			
Some gene mutations change only one triplet code. Due to the degenerate nature of the genetic code, not all such mutations result in a change to the encoded amino acid			
Some gene mutations change the nature of all base triplets downstream from the mutation, ie result in a frame shift			
Be able to relate the nature of a gene mutation to its effect on the encoded polypeptide			

GENE EXPRESSION IS CONTROLLED BY A NUMBER OF FEATURES			
Most of a cell's DNA is not translated			
Totipotent cells can divide and produce any type of body cell			
During development, totipotent cells translate only part of their DNA, resulting in cell specialisation			
Totipotent cells occur only for a limited time in early mammalian embryos			
Pluripotent cells are found in embryos; multipotent and unipotent cells are found in mature mammals and can divide to form a limited number of different cell types			
Pluripotent stem cells can divide in unlimited numbers and can be used in treating human disorders			
Unipotent cells, exemplified by the formation of cardiomyocytes			

• Induced pluripotent stem cells (iPS cells) can be produced from adult somatic cells using appropriate protein transcription factors			
Be able to evaluate the use of stem cells in treating human disorders			
Regulation of transcription and translation			
In eukaryotes, transcription of target genes can be stimulated or inhibited when specific transcriptional factors move from the cytoplasm into the nucleus. The role of the steroid hormone, oestrogen, in initiating transcription			
Epigenetic control of gene expression in eukaryotes			
Epigenetics involves heritable changes in gene function, without changes to the base sequence of DNA. These changes are caused by changes in the environment that inhibit transcription by:			
• increased methylation of the DNA, or:			
• decreased acetylation of associated histones			
The relevance of epigenetics on the development and treatment of disease, especially cancer			
In eukaryotes and some prokaryotes, translation of the mRNA produced from target genes can be inhibited by RNA interference (RNAi)			
Be able to interpret data provided from investigations into gene expression			
Be able to evaluate appropriate data for the relative influences of genetic and environmental factors on phenotype			
Gene expression and cancer			
The main characteristics of benign and malignant tumours			
The role of the following in the development of tumours:			
• tumour suppressor genes and oncogenes			
• abnormal methylation of tumour suppressor genes and oncogenes			
• increased oestrogen concentrations in the development of some breast cancers			
Be able to evaluate evidence showing correlations between genetic and environmental factors and various forms of cancer			
Be able to interpret information relating to the way in which an understanding of the roles of oncogenes and tumour suppressor genes could be used in the prevention, treatment and cure of cancer			

USING GENOME PROJECTS			
Sequencing projects have read the genomes of a wide range of organisms, including humans			
Determining the genome of simpler organisms allows the sequences of the proteins that derive from the genetic code (the proteome) of the organism to be determined. This may have many applications, including the identification of potential antigens for use in vaccine production			

In more complex organisms, the presence of non-coding DNA and of regulatory genes means that knowledge of the genome cannot easily be translated into the proteome			
Sequencing methods are continuously updated and have become automated			

GENE TECHNOLOGIES ALLOW THE STUDY OF ALTERATION OF GENE FUNCTION ALLOWING A BETTER UNDERSTANDING OF ORGANISM FUNCTION AND THE DESIGN OF NEW INDUSTRIAL AND MEDICAL PROCESSES			
Recombinant DNA technology			
Recombinant DNA technology involves the transfer of fragments of DNA from one organism, or species, to another. Since the genetic code is universal, as are transcription and translation mechanisms, the transferred DNA can be translated within cells of the recipient (transgenic) organism			
Fragments of DNA can be produced by several methods, including:			
• conversion of mRNA to complementary DNA (cDNA), using reverse transcriptase			
• using restriction enzymes to cut a fragment containing the desired gene from DNA			
• creating the gene in a 'gene machine'			
Fragments of DNA can be amplified by in vitro and in vivo techniques			
The principles of the polymerase chain reaction (PCR) as an <i>in vitro</i> method to amplify DNA fragments			
The culture of transformed host cells as an in vivo method to amplify DNA fragments			
• The addition of promoter and terminator regions to the fragments of DNA			
• The use of restriction endonucleases and ligases to insert fragments of DNA into vectors. Transformation of host cells using these vectors			
• The use of marker genes to detect genetically modified (GM) cells or organisms			
Students will not be required to recall specific marker genes in a written paper			
Be able to interpret information relating to the use of recombinant DNA technology			
Be able to evaluate the ethical, financial and social issues associated with the use and ownership of recombinant DNA technology in agriculture, in industry and in medicine			
Be able to balance the humanitarian aspects of recombinant DNA technology with the opposition from environmentalists and anti-globalisation activists			
Be able to relate recombinant DNA technology to gene therapy			
Differences in DNA between individuals of the same species can be exploited for identification and diagnosis of heritable conditions			
The use of labelled DNA probes and DNA hybridisation to locate specific alleles of genes			

The use of labelled DNA probes that can be used to screen patients for heritable conditions, drug responses or health risks			
The use of this information in genetic counselling and personalised medicine			
Be able to evaluate information relating to screening individuals for genetically determined conditions and drug responses			
Genetic fingerprinting			
An organism's genome contains many variable number tandem repeats (VNTRs). The probability of two individuals having the same VNTRs is very low			
The technique of genetic fingerprinting in analysing DNA fragments that have been cloned by PCR, and its use in determining genetic relationships and in determining the genetic variability within a population			
The use of genetic fingerprinting in the fields of forensic science, medical diagnosis, animal and plant breeding			
Be able to explain the biological principles that underpin genetic fingerprinting techniques			
Be able to interpret data showing the results of gel electrophoresis to separate DNA fragments			
Be able to explain why scientists might use genetic fingerprinting in the fields of forensic science, medical diagnosis, animal and plant breeding			



Induction tasks


**To be completed and handed in to
your teacher in September**

Name: _____


Cells are a bit more complicated...

Task - Draw and label a GCSE level diagram of a eukaryotic animal cell, then draw a labelled diagram of this cell type that is A level quality. Remember that your exam board is AQA, so pay close attention to what detail is expected of you for this course.

GCSE diagram



'A' Level diagram

A large, empty rectangular box with rounded corners, intended for drawing an 'A' level diagram. The box is white and occupies the majority of the page below the title.

Biological molecules underpin everything that we study at A level, so you need to start to develop a working knowledge of the main examples.

monosaccharide	deoxyribose	condensation reaction	R group	monomer
hydrogen bonding	triglyceride	amine group	polypeptide chain	nitrogenous base
CHO	glycosidic bond	sucrose	polynucleotide	saturated
polysaccharide	isomer	CHON	inorganic phosphate	hydrolysis
beta-pleated sheet	unsaturated	glycerol	primary structure	carboxyl group
CHONP	complementary	nucleic acid	β -glucose	disaccharide
maltose	secondary structure	phosphodiester bond	CHO	amylose
cellulose	disulfide bridge	quaternary structure	hydrophobic	pentose
hexose	water	glycogen	ionic bonding	microfibril
phospholipid	hydrophilic	fatty acid	polar	nucleotide
ribose	peptide bond	alpha helix	ester bond	Starch
fructose	amylopectin	tertiary structure	base pairing	covalent bonding

Step 1 – Become familiar with Miss Estruch, a firm favourite of past KS5 years by watching this video on biological molecules ([19\) A level Biological Molecules - Learn the ENTIRE topic in this video. AQA A level Biology Revision - YouTube](#)). It is a long-ish video, and does contain additional content, but that will never be harmful.

Step 2 – sort the terms in the table above into the following categories: Lipids, Proteins, Nucleic acids and Carbohydrates use the resources below for support.



Science Hive

[Biological Molecules \(A Level\) — the science hive](#)



Save my exams

[AQA A Level Biology Revision Notes 2017 | Save My Exams](#)



PMT

[AQA A-level Biology Revision - PMT](#)



Free science lessons

[A-Level Biology | freesciencelessons](#)

Step 3 – Write a minimum of two sides of A4 on each of the biological molecules, you must include relevant diagrams. (Complete this task on lined paper)

Writing like and A level biologist

You have learnt a fair quantity of the vocabulary that you will need for being successful in A level biology, however, there is no getting away from the fact that as A level biologists your vocabulary needs to become even more sophisticated and nuanced.

In this task for every GCSE key term provided you must find the terms that you will be using instead to enhance your writing and make you sound like an A level biologist.

GCSE term – White blood cells	GCSE understanding/knowledge of this key term
A level term(s) that I will be using instead...	
Useful detail I discover whilst searching for the terminology...	

GCSE term – Lock and key model	GCSE understanding/knowledge of this key term
A level term(s) that I will be using instead...	
Useful detail I discover whilst searching for the terminology...	

GCSE term – Sugar	GCSE understanding/knowledge of this key term
A level term(s) that I will be using instead...	
Useful detail I discover whilst searching for the terminology...	

GCSE term – Membrane	GCSE understanding/knowledge of this key term
A level term(s) that I will be using instead...	
Useful detail I discover whilst searching for the terminology...	

GCSE term – Lipid	GCSE understanding/knowledge of this key term
A level term(s) that I will be using instead...	
Useful detail I discover whilst searching for the terminology...	

GCSE term – Energy	GCSE understanding/knowledge of this key term
A level term(s) that I will be using instead...	
Useful detail I discover whilst searching for the terminology...	

GCSE term – Water level	GCSE understanding/knowledge of this key term
A level term(s) that I will be using instead...	
Useful detail I discover whilst searching for the terminology...	

1 Numbers and units

1.1 Units and prefixes

A key criterion for success in biological maths lies in the use of correct units and the management of numbers. The units scientists use are from the *Système Internationale* – the SI units. In biology, the most commonly used SI base units are metre (m), kilogram (kg), second (s), and mole (mol). Biologists also use SI derived units, such as square metre (m²), cubic metre (m³), degree Celsius (°C), and litre (l).

To accommodate the huge range of dimensions in our measurements they may be further modified using appropriate prefixes. For example, one thousandth of a second is a millisecond (ms). Some of these prefixes are illustrated in the table below.

Multiplication factor	Prefix	Symbol
10 ⁹	giga	G
10 ⁶	mega	M
10 ³	kilo	k
10 ⁻²	centi	c
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n

Practice questions

- A burger contains 4 500 000 J of energy.
Write this in: **a** kilojoules **b** megajoules.
- HIV is a virus with a diameter of between 9.0×10^{-8} m and 1.20×10^{-7} m.
Write this range in nanometres.

1.2 Powers and indices

Ten squared = $10 \times 10 = 100$ and can be written as 10^2 . This is also called 'ten to the power of 2'. Ten cubed is 'ten to the power of three' and can be written as $10^3 = 1000$. The power is also called the index. Fractions have negative indices:

one tenth = $10^{-1} = 1/10 = 0.1$

one hundredth = $10^{-2} = 1/100 = 0.01$

Any number to the power of 0 is equal to 1, for example, $29^0 = 1$.

If the index is 1, the value is unchanged, for example, $17^1 = 17$. When multiplying powers of ten, you must *add* the indices. So, $100 \times 1000 = 100\,000$ is the same as $10^2 \times 10^3 = 10^{2+3} = 10^5$

When dividing powers of ten, you must *subtract* the indices. So, $100/1000 = 1/10 = 10^{-1}$ is the same as $10^2/10^3 = 10^{2-3} = 10^{-1}$

But you can only do this when the numbers with the indices are the same. So, $10^2 \times 2^3 = 100 \times 8 = 800$

And you can't do this when adding or subtracting. $10^2 + 10^3 = 100 + 1000 = 1100$

$10^2 - 10^3 = 100 - 1000 = -900$

Remember: You can only add and subtract the indices when you are multiplying or dividing the numbers, not adding, or subtracting them.

Practice questions

3 Calculate the following values. Give your answers using indices.

- a $10^8 \times 10^3$ b $10^7 \times 10^2 \times 10^3$
c $10^3 + 10^3$ d $10^2 - 10^{-2}$

4 Calculate the following values. Give your answers with and without using indices.

- a $10^5 \div 10^4$ b $10^3 \div 10^6$
c $10^2 \div 10^{-4}$ d $100^2 \div 10^2$

1.3 Converting units

When doing calculations, it is important to express your answer using sensible numbers. For example, an answer of $6230\,\mu\text{m}$ would have been more meaningful expressed as $6.2\,\text{mm}$.

If you convert between units and round numbers properly, it allows quoted measurements to be understood within the scale of the observations.

To convert $488\,889\,\text{m}$ into km :

A kilo is 10^3 so you need to divide by this number or move the decimal point three places to the left.

$488\,889 \div 10^3 = 488.889\,\text{km}$

However, suppose you are converting from mm to km : you need to go from 10^3 to 10^{-3} , or move the decimal point six places to the left.

$333\,\text{mm}$ is $0.000\,333\,\text{km}$

Alternatively, if you want to convert from $333\,\text{mm}$ to nm , you will have to go from 10^{-9} to 10^{-3} , or move the decimal point six places to the right.

$333\,\text{mm}$ is $333\,000\,000\,\text{nm}$

Practice questions

- 5 Calculate the following conversions:
a 0.004 m into mm b 130 000 ms into s
c 31.3 ml into μl d 104 ng into mg
- 6 Give the following values in a different unit so they make more sense to the reader.
Choose the final units yourself. (Hint: make the final number as close in magnitude to zero as you can. For example, you would convert 1000 m into 1 km.)
a 0.000 057 m b 8 600 000 μl c 68 000 ms d 0.009 cm

2 Decimals, standard form, and significant figures

2.1 Decimal numbers

A decimal number has a decimal point. Each figure *before* the point is a whole number, and the figures *after* the point represent fractions.

The number of decimal places is the number of figures *after* the decimal point. For example, the number 47.38 has 2 decimal places, and 47.380 is the same number to 3 decimal places.

In science, you must write your answer to a sensible number of decimal places.

Practice questions

- 1 New antibiotics are being tested. A student calculates the area of clear zones in Petri dishes in which the antibiotics have been used. List these in order from smallest to largest.
0.0214 cm^2 0.03 cm^2 0.0218 cm^2 0.034 cm^2
- 2 A student measures the heights of a number of different plants. List these in order from smallest to largest.
22.003 cm 22.25 cm 12.901 cm 12.03 cm 22 cm

2.2 Standard form

Sometimes biologists need to work with numbers that are very small, such as dimensions of organelles, or very large, such as populations of bacteria. In such cases, the use of scientific notation or standard form is very useful, because it allows the numbers to be written easily.

Standard form is expressing numbers in powers of ten, for example, 1.5×10^7 microorganisms.

Look at this worked example. The number of cells in the human body is approximately 37 200 000 000 000. To write this in standard form, follow these steps:

Step 1: Write down the smallest number between 1 and 10 that can be derived from the number to be converted. In this case it would be 3.72

Step 2: Write the number of times the decimal place will have to shift to expand this to the original number as powers of ten. On paper this can be done by hopping the decimal over each number like this: until the end of the number is reached.

In this example that requires 13 shifts, so the standard form should be written as 3.72×10^{13} .

For very small numbers the same rules apply, except that the decimal point has to hop backwards. For example, 0.000 000 45 would be written as 4.5×10^{-7} .

Practice questions

3 Change the following values to standard form.

- a 3060 kJ b 140 000 kg c 0.000 18 m d 0.000 004 m

4 Give the following numbers in standard form.

- a 100 b 10 000 c 0.01 d 21 000 000

5 Give the following as decimals.

- a 10^6 b 4.7×10^9 c 1.2×10^{12} d 7.96×10^{-4}

2.3 Significant figures

When you use a calculator to work out a numerical answer, you know that this often results in a large number of decimal places and, in most cases, the final few digits are 'not significant'. It is important to record your data and your answers to calculations to a reasonable number of significant figures. Too many and your answer is claiming an accuracy that it does not have, too few and you are not showing the precision and care required in scientific analysis.

Numbers to 3 significant figures (3 s.f.):

7.88 25.4 741

Bigger and smaller numbers with 3 significant figures:

0.000 147 0.0147 0.245 39 400 96 200 000 (notice that the zeros before the figures and after the figures are *not* significant – they just show you how large the number is by the position of the decimal point).

Numbers to 3 significant figures where the zeros *are* significant:

207 4050 1.01 (any zeros between the other significant figures *are* significant).

Standard form numbers with 3 significant figures:

9.42×10^{-5} 1.56×10^8

If the value you wanted to write to 3.s.f. was 590, then to show the zero was significant you would have to write:

590 (to 3.s.f.) or 5.90×10^2

Remember: For calculations, use the same number of figures as the data in the question with the lowest number of significant figures. It is not possible for the answer to be more accurate than the data in the question.

Practice question

6 Write the following numbers to **i** 2 s.f. and **ii** 3 s.f.

a 7644 g

b 27.54 m

c 4.3333 g

d $5.995 \times 10^2 \text{ cm}^3$

7 The average mass of oxygen produced by an oak tree is 11800 g per year. Give this mass in standard form and quote your answer to 2 significant figures.